

Microbial pathogens in ticks, rodents and a shrew in northern Gyeonggi-do near the DMZ, Korea

Joon-Seok Chae^{1,*}, Do-Hyeon Yu², Smriti Shringi², Terry A. Klein³, Heung-Chul Kim⁴, Sung-Tae Chong⁴, In-Yong Lee⁵, Janet Foley⁶

¹Department of Internal Medicine, College of Veterinary Medicine, Seoul National University, Seoul 151-742, Korea

²College of Veterinary Medicine, Chonbuk National University, Jeonju 561-756, Korea

³Force Health Protection, 18th Medical Command, Unit #15281, Box 754, APO AP 96205-5281, USA

⁴5th Medical Detachment, 168th Multifunctional Medical Battalion, 18th Medical Command, Unit #15247, APO AP 96205-5247, USA

⁵Department of Environmental Medical Biology, College of Medicine, Yonsei University, Seoul 120-749, Korea

⁶Center for Vector-Borne Diseases, School of Veterinary Medicine, University of California, Davis, CA 95616, USA

A total of 1,618 ticks [420 individual (adults) and pooled (larvae and nymphs) samples], 369 rodents (*Apodemus agrarius*, *Rattus norvegicus*, *Tscherskia triton*, *Mus musculus*, and *Myodes regulus*), and 34 shrews (*Crocidura lasiura*) that were collected in northern Gyeonggi-do near the Demilitarized Zone (DMZ) of Korea during 2004-2005, were assayed by PCR for selected zoonotic pathogens. From a total of 420 individual and pooled tick DNA samples, *Anaplasma (A.) phagocytophilum* (16), *A. platys* (16), *Ehrlichia (E.) chaffeensis* (63), *Borrelia burgdorferi* (16), and *Rickettsia* spp. (198) were detected using species-specific PCR assays. Out of 403 spleens from rodents and shrews, *A. phagocytophilum* (20), *A. platys* (34), *E. chaffeensis* (127), and *Bartonella* spp. (24) were detected with species-specific PCR assays. These results suggest that fevers of unknown causes in humans and animals in Korea should be evaluated for infections by these vector-borne microbial pathogens.

Keywords: Bartonella, Borrelia, Rickettsia, rodents, *Crocidura lasiura*, tick-borne pathogens

Introduction

Korea is a northeast Asian peninsular country with four clearly demarked seasons. Seventy percent of the land area is mountainous, with interspersed fertile river valleys. Ticks are commonly collected during the early spring through late autumn, while few ticks are collected

during the cold winter season. Many wild animals inhabit the Demilitarized Zone (DMZ) and the area adjacent to it, and these animals are hosts to ticks and serve as reservoirs for tick-borne pathogens [17]. The Korean and US military have numerous small to large training sites near the DMZ where large populations of small mammals (rodents and insectivores) and occasional deer, wild pigs, and other small mammals are found [32]. Additionally, tourist activity is expected to increase in the area in the near future, which may increase the risk of human exposure to ticks and the pathogens they harbor [5,16].

Ectoparasites (e.g. ticks and fleas) are vectors of a number of pathogens that are important to humans and also veterinary practice. Ticks are harmful ectoparasites that directly or indirectly cause a variety of disease states in their host. Ticks are known vectors of protozoa, rickettsiae, bacteria, and viruses, that may cause serious and life-threatening illnesses in human. Screening ticks for disease-causing pathogens using molecular epidemiological tools provides useful data on the distribution and prevalence of tick-borne pathogens. Moreover, with increases in the mean global annual temperatures of 1°C since the 1880s [10], it is predicted that the temperate Korean climate may be altered to a subtropical climate. These environmental changes may potentially alter the distribution of wild animals and the arthropod vectors and the pathogens they transmit. Tick-borne encephalitis was previously thought to not exist in Korea, but recent evidence from molecular testing of ticks and rodents suggests that it is present in Korea [19]. Many of the pathogenic agents transmitted by ticks, including *Ehrlichia* spp., *Anaplasma* spp., *Borrelia* spp., *Bartonella* spp., and *Rickettsia* spp., are known to be human and animal pathogens worldwide [8,20,29].

*Corresponding author

Tel: +82-2-880-1279; Fax: +82-02-880-1213

E-mail: jschae@snu.ac.kr

Recent seroepidemiological findings documented the presence of human monocytic ehrlichiosis and human granulocytic anaplasmosis in Korea [11,26]. Molecular evidence of *Ehrlichia* and *Anaplasma* spp. was identified in ticks collected from animals and grass vegetation in Korea [17,21]. Additionally, a spotted fever group Rickettsia, similar to *Rickettsia* (*R.*) *japonica*, was identified in *Haemaphysalis* (*H.*) *longicornis* ticks by PCR, and antibodies to these organisms were detected in human patients with acute febrile illness [14].

The United States Forces Korea rodent- and tick-borne disease surveillance program was initiated to provide ecological and epidemiological information on potential risks of infection for personal who occupy or train in various environments near the DMZ. This is especially important when considering recent serological evidence that confirmed the presence of *Ehrlichia* (*E.*) *chaffeensis* and *Anaplasma* (*A.*) *phagocytophilum* [11,26].

The purpose for this study was to identify vector-borne pathogens in ticks, rodents and shrews in order to provide more accurate risk assessment of tick-borne pathogens that may affect human and animal health in Korea.

Materials and Methods

Study sites

Ticks were collected in the field by dragging and flagging grass vegetation and forested ground cover (fallen leaves, clumps of grasses and scattered shrubs). Ticks also were removed from various wild rodents (*Apodemus agrarius*, *Rattus norvegicus*, *Tscherskia triton*, *Mus musculus*, and *Myodes regulus*) and a shrew (*Crocidura lasiura*) that were live-trapped at US military installations and training sites in northern Gyeonggi-do near the DMZ (Fig. 1).

Tick collections

During March through September 2004, a total of 1,618 ticks were collected from grass vegetation and forest leaf litter (933 ticks) and wild rodents (685 ticks) at 17 sites (Fig. 1). Based on microscopic examination, ticks were identified to species and developmental stage characterized. Adult ticks were stored and assayed individually, while the nymphs and larvae were pooled (1-6 and 1-30 ticks per pool, respectively) into 420 sample pools (62 from wild rodents and 358 from grass vegetation and forest leaf litter) and stored at -70°C until they were assayed.

Tissue samples

A total of 403 small mammals (369 wild rodents and 34 shrews) belonging to six species, six different genera, and two families were live captured at US military installations and training sites in northern Gyeonggi-do near the DMZ in Korea from August 2004 through June 2005 using Sherman traps (3" × 5" × 9" folding traps; H.B. Sherman

Traps, USA). The live-caught rodents and shrews were transported to Korea University where they were euthanized in accordance with the Korea University animal use protocol, their abdominal cavities opened aseptically, and spleen samples collected and stored individually at -70°C until assayed.

DNA extraction

DNA was extracted from pools of larvae, nymphs and individual adult ticks. A total of 747 and 174 nymphs were collected by tick drag/flag and from rodents and a shrew, respectively, and these were placed in 215 pools according to collection site, while DNA was extracted from 186 individual adult ticks (76 males and 110 females) and 19 pools of larvae with using DNeasy tissue kits (Qiagen, Germany) (Table 1). Individual ticks and pools of ticks were mechanically homogenized using sterile scissors and a manual homogenizer (General Biosystem, Korea). DNA extraction was performed using DNeasy tissue kits (Qiagen, Germany) in accordance with instructions provided by the manufacturer.

Detection of tick-borne pathogens by PCR

Purified DNA was used for the detection of tick-borne pathogens using conventional and nested PCR [16]. PCR assays using genomic DNAs and species-specific primers, as previously described, were used to identify selected zoonotic pathogens [18].

Nested PCR: The nested PCR technique was used for the detection of *A. phagocytophilum* by amplifying a 926 bp fragment of *A. phagocytophilum*-specific 16S rRNA gene in a total volume of 25 µl as previously described [4]. Species-specific primers for *A. platys*, *E. chaffeensis*, *E. ewingii*, and *E. canis* were used in the nested PCR assays [23,24]. The primers ECC and ECB were used to amplify




Fig. 1. Collection sites were conducted in northern Gyeonggi-do near the Demilitarized Zone of Korea. The small black squares indicate sample collection sites.

all *Ehrlichia* spp. [6,7]. The primers EPLAT5 and EPLAT3 were used for *A. platys*-specific amplification [22], the primers HE1 and HE3 were used for *E. chaffeensis*-specific amplification [3], the primers EE52 and HE3 were used for *E. ewingii*-specific amplification [23], and the primers ECAN5 and HE3 were used for *E. canis*-specific amplification [23].

Conventional PCR: Identification of *Bartonella* spp., *E. muris*, *Borrelia* (*B.*) *burgdorferi*, *R. rickettsii*, and *R. japonica* was performed using conventional PCR with the species-specific primers [9,30,33]. The citrate synthase gene (*gltA*) was selected for the identification of *E. muris* [14]. The primers BhCS (781p) and BhCS (1137n) were used for *Bartonella* spp. amplification [24]. The *gltA* gene was used for the identification of *Bartonella* spp. The *ospC* gene was selected for the identification of *B. burgdorferi*. A pair of synthesized oligonucleotide primers derived from the gene sequence encoding the 190 kDa antigen of *R. rickettsii*, Rr190.70p and Rr190.602n, as described by Regnery *et al.* [30], was used for the PCR amplification of *R. rickettsii* DNA. Species-specific primers, Rj10 and Rj5, were used for the *R. japonica* 17 kDa antigen gene fragment [9]. PCR reactions were performed using 50-100 ng of template DNA, a species-specific primer set, and the PCR mixture. The PCR products were electrophoresed in 1% agarose gel; they were then stained with ethidium bromide and photographed using a still video documentation system (Gel Doc 2000; BioRad, USA).

Isolation of *Bartonella* sp.

Small mammal spleens were collected in 2 ml tubes and maintained on dry ice for transportation and subsequently used for culture isolation. The spleens were homogenized and then plated on fresh chocolate agar and allowed to incubate in 5% CO₂ at 35°C for up to 4 weeks. The single colonies that grew were scraped for identification of *Bartonella* spp. The isolates were then confirmed as *Bartonella* spp. by PCR and DNA sequencing. Culture isolates were stored at -70°C in frozen medium [a total of 100 ml; M199 tissue culture medium with glutamine and Earle's salts (GIBCO, USA), 1 ml of ×100 glutamine (GIBCO, USA), 1 ml of ×100 sodium pyruvate (GIBCO, USA), 20% bovine fetal calf serum (heat inactivated), and 3 ml sodium bicarbonate (7.5% solution) (GIBCO, USA), 10% DMSO, pH: 7.1-7.4] for later use.

Results

A total of 1,618 ticks from two genera and three species [570 *H. longicornis*, 306 *H. flava* and 742 *Ixodes* (*I.*) *nipponensis*] was collected from grass vegetation and forest leaf litter (933 ticks) and small mammals (685 ticks) from 2004 to 2005 near or at US military installations and

training sites in northern Gyeonggi-do near the DMZ, Korea (Fig. 1, Table 1). *H. longicornis* ticks were the most frequently collected species from the grass fields. Except for one *H. flava*, all ticks taken from captured small mammals were *I. nipponensis* larvae and nymphs (Table 1).

Species-specific PCR assays were performed using DNA samples from 420 individuals and pools of ticks, and DNA samples from spleens of 403 small mammals. Five of the ten tick-borne pathogens examined in this study were detected in ticks [*A. phagocytophilum* (16, 1.0%), *A. platys* (16, 1.0%), *E. chaffeensis* (63, 3.9%), *B. burgdorferi* (16, 1.0%), and *Rickettsia* spp. (198, 12.2%)] (Table 1). At least fifty-one ticks had a mixed infection with two pathogens: *E. chaffeensis* and *Rickettsia* spp. (32 samples), *A. phagocytophilum* and *E. chaffeensis* (3 samples), *A. phagocytophilum* and *Rickettsia* spp. (4 samples), *Rickettsia* spp. and *A. platys* (3 samples), *B. burgdorferi* and *Rickettsia* spp. (6 samples), *E. chaffeensis* and *B. burgdorferi* (2 samples), and *A. phagocytophilum* and *B. burgdorferi* (1 sample) (Table 2). At least eight ticks had mixed infections with three pathogens: *A. platys*, *E. chaffeensis* and *Rickettsia* spp. (5 samples), *B. burgdorferi*, *Rickettsia* spp. and *A. phagocytophilum* (2 samples), and *A. phagocytophilum*, *A. platys* and *Rickettsia* spp. (1 sample) (Table 2).

A total of 403 small mammals were collected from US military installations and training sites in northern Gyeonggi-do near the DMZ, and these included five rodents, *Apodemus agrarius* (358), *Rattus norvegicus* (6), *Tscherskia triton* (2), *Mus musculus* (2), *Myodes regulus* (1) and a shrew, *Crocidura lasiura* (34) (Table 3). Four of the ten tick-borne pathogens examined in this study were detected by PCR in the small mammals [*A. phagocytophilum* (20, 5.0%), *A. platys* (34, 8.4%), *E. chaffeensis* (127, 31.5%) and *Bartonella* spp. (24, 6.0%)] (Table 3). *Apodemus agrarius* was PCR positive for *A. phagocytophilum*, *A. platys*, *E. chaffeensis* and *Bartonella* spp., while *Mus musculus* was only positive for *E. chaffeensis*. *Crocidura lasiura* was positive only for *A. platys* and *E. chaffeensis* (Table 3).

A total of 376 small mammals had single infections with rickettsial pathogens, while 26 *Apodemus agrarius* had mixed infections of two (23 samples), or three (3 samples) pathogens and a single *Crocidura lasiura* was positive for two pathogens (Table 4).

The frozen and homogenized samples of spleens of *Apodemus agrarius* were cultured and grew as a non-hemolytic gram-negative organism after 14 days, with only a few small white colonies. PCR amplification from the 10 isolates using *gltA* primers produced a 356 bp fragment and sequencing results were strongly suggestive of *Bartonella elizabethae* by phylogenetic analysis [17].

Discussion

An analysis of ticks and small mammal tissues demonstrated a high rate of infection of tick-borne pathogens in northern Gyeonggi-do near the DMZ. Most *Ehrlichia* and *Anaplasma* spp. tick-borne infections occur in *Ixodes* spp. in the US and Europe [1,31]. In Asia, *Ehrlichia* spp. was previously identified from *Haemaphysalis* spp. as well as *Ixodes* spp. [13,17,18]. *H. longicornis* are widespread throughout Korea, and especially around the pastures for grazing cattle or where deer congregate.

I. nipponensis are two-host ticks with larvae and nymphs found on rodents and a shrew. Infection rates of *Rickettsia* spp. (56.5%) and *B. burgdorferi* (25.8%) were relatively high among the selected rodents and a shrew tested. Ticks collected from grass vegetation and forest leaf litter were negative for *B. burgdorferi*, which may be a result of the small sample size of *I. nipponensis* from the "collected vegetation". In experimentally infected mice, *B. burgdorferi* DNA can be detected from the foot and lymph nodes by PCR until 55 days post-inoculation [25]. In that study, *B. burgdorferi* DNA was detected from the spleen tissues 15 days post inoculation, but not at 55 days post inoculation. Persistent infections have also been reported in the skin, blood, CSF and synovial fluid of human patients [2,25]. In the present study, *B. burgdorferi* DNA was not detected from the spleens of rodents and a shrew or the ticks, but was identified from the *I. nipponensis* removed from the small mammals. This suggests that wild rodents are a natural reservoir of *B. burgdorferi* in Korea, with *I. nipponensis* as an important vector for the larger animal hosts.

In this study, there was a very high prevalence of *Rickettsia* spp. in *H. longicornis*, *H. flava* and *I. nipponensis* ticks, but not in rodents and a shrew. Our previous studies during 2001 through 2003 detected *Rickettsia* spp. only from *H. longicornis* and *Apodemus agrarius* [18]. The PCR primer set in the previous studies targeted the *R. rickettsii* *rOmpA* gene [30], and we were able to sequence the amplicons. The resultant phylogenetic tree showed that Korean rickettsias were closely related to the *Rickettsia* spp. strain FUJ98 in China [18].

Additionally, these results showed that only one *Ixodes* spp. tick collected from vegetation was found infected with *A. phagocytophilum* (0.1%) [18]. In the present study, the *A. phagocytophilum* infection rate observed in rodents and a shrew tissues (5.6%) was similar to the rate of infection for *I. nipponensis* ticks collected from rodents and a shrew (5.2%), while only 1.8% of *I. nipponensis* collected from vegetation were positive for *A. phagocytophilum*.

Specific DNA of *E. canis*, *E. ewingii*, *E. muris* and *R. japonica* was not amplified in this study. There have been previous reports of the spotted fever group rickettsiosis, including *R. japonicus*, in Korean patients and ticks

[15,28].

Our results demonstrate that ticks and rodents and a shrew captured near the DMZ of Korea were infected with *Anaplasma*, *Ehrlichia*, *Bartonella*, *Borrelia*, and *Rickettsia* spp. Although infections with *Ehrlichia* and *Anaplasma* spp. have generally been considered to be observed only in a defined range of hosts, including rodents and some large mammals, our studies suggest that several *Ehrlichia* and *Anaplasma* spp. can be transmitted to a variety of hosts in nature. Therefore, additional efforts to define the spectrum of host susceptibility in domestic and wild animals are needed.

H. longicornis, *H. flava* and *I. nipponensis* should be considered as potential vectors of *A. phagocytophilum*, *A. platys*, *E. chaffeensis* and *Rickettsia* spp., while *Apodemus agrarius*, *Crocidura lasiura* and *Mus musculus* may be reservoir hosts of selected tick-borne pathogens in Korea.

Until now, there have not been reports of clinical cases for *A. phagocytophilum*, *E. chaffeensis* and *B. elizabethae* in humans and animals in the Korea, as compared with the numerous reports throughout the world. For some diseases, such as rabies and malaria, there have been reported outbreaks along the DMZ [12,27]. Therefore, in the future, it will become important to perform surveillance for pathogens, including *Anaplasma*, *Ehrlichia*, *Bartonella*, *Borrelia*, and *Rickettsia* spp., in vectors and wild animals, as well as in civilian and military populations that reside or train near the DMZ. It is imperative to continue the efforts to identify additional tick-borne pathogens to further disclose the extent and possible public health significance of these agents.

Acknowledgments

Funding for portions of this work was provided by the US Department of Defense Global Emerging Infections Surveillance and Response System, Silver Spring, MD, the Armed Forces Medical Intelligence Center, Ft Detrick, MD. Dr. Joon-seok Chae received funding from the LG Yeonam Foundation.

References

1. Adelson ME, Rao RVS, Tilton RC, Cabets K, Eskow E, Fein L, Occi JL, Mordechai E. Prevalence of *Borrelia burgdorferi*, *Bartonella* spp. *Babesia microti*, and *Anaplasma phagocytophilum* in *Ixodes scapularis* ticks collected in northern New Jersey. *J Clin Microbiol* 2004, **42**, 2799-2801.
2. Agüero-Rosenfeld ME, Wang G, Schwartz I, Wormser GP. Diagnosis of Lyme borreliosis. *Clin Microbiol Rev* 2005, **18**, 484-509.
3. Anderson BE, Sumner JW, Dawson JE, Tzianabos T, Greene CR, Olson JG, Fishbein DB, Olsen-Rasmussen M, Holloway BP, George EH, Azad AF. Detection of the etiologic agent of human ehrlichiosis by polymerase chain

- reaction. *J Clin Microbiol* 1992, **30**, 775-780.
4. **Barlough JE, Madigan JE, Derock E, Bigornia L.** Nested polymerase chain reaction for detection of *Ehrlichia equi* genomic DNA in horses and ticks (*Ixodes pacificus*). *Vet Parasitol* 1996, **63**, 319-329.
 5. **Chae JS, Kim CM, Kim EH, Hur EJ, Klein TA, Kang TK, Lee HC, Song JW.** Molecular epidemiological study for tick-borne disease (*Ehrlichia* and *Anaplasma* spp.) surveillance at selected U.S. military training sites/installations in Korea. *Ann N Y Acad Sci* 2003, **990**, 118-125.
 6. **Dawson JE, Biggie KL, Warner CK, Cookson K, Jenkins S, Levine JF, Olson JG.** Polymerase chain reaction evidence of *Ehrlichia chaffeensis*, an etiologic agent of human ehrlichiosis, in dogs from southeast Virginia. *Am J Vet Res* 1996, **57**, 1175-1179.
 7. **Dawson JE, Stallknecht DE, Howerth EW, Warner C, Biggie K, Davidson WR, Lockhart JM, Nettles VF, Olson JG, Childs JE.** Susceptibility of white-tailed deer (*Odocoileus virginianus*) to infection with *Ehrlichia chaffeensis*, the etiologic agent of human ehrlichiosis. *J Clin Microbiol* 1994, **32**, 2725-2728.
 8. **Fritz CL, Kjemtrup AM.** Lyme borreliosis. *J Am Vet Med Assoc* 2003, **223**, 1261-1270.
 9. **Furuya Y, Katayama T, Yoshida Y, Kaiho I.** Specific amplification of *Rickettsia japonica* DNA from clinical specimens by PCR. *J Clin Microbiol* 1995, **33**, 487-489.
 10. **Hansen J, Ruedy R, Glascoe J, Sato N.** GISS analysis of surface temperature change. *J Geophys Res* 1999, **104**, 30997-31022.
 11. **Heo EJ, Park JH, Koo JR, Park MS, Park MY, Dumler JS, Chae JS.** Serologic and molecular detection of *Ehrlichia chaffeensis* and *Anaplasma phagocytophila* (Human granulocytic ehrlichiosis agent) in Korean patients. *J Clin Microbiol* 2002, **40**, 3082-3085.
 12. **Hyun BH, Lee KK, Kim IJ, Lee KW, Park HJ, Lee OS, An SH, Lee JB.** Molecular epidemiology of rabies virus isolates from South Korea. *Virus Res* 2005, **114**, 113-125.
 13. **Inokuma H, Beppu T, Okuda M, Shimada Y, Sakata Y.** Detection of ehrlichial DNA in *Haemaphysalis* ticks recovered from dogs in Japan that is closely related to a novel *Ehrlichia* sp. found in cattle ticks from Tibet, Thailand, and Africa. *J Clin Microbiol* 2004, **42**, 1353-1355.
 14. **Inokuma H, Brouqui P, Drancourt M, Raoult D.** Citrate synthase gene sequence: a new tool for phylogenetic analysis and identification of *Ehrlichia*. *J Clin Microbiol* 2001, **39**, 3031-3039.
 15. **Jang WJ, Kim JH, Choi YJ, Jung KD, Kim YG, Lee SH, Choi MS, Kim IS, Walker DH, Park KH.** First serologic evidence of human spotted fever group rickettsiosis in Korea. *J Clin Microbiol* 2004, **42**, 2310-2313.
 16. **Kim CM, Kim JY, Yi YH, Lee MJ, Cho MR, Shah DH, Klein TA, Kim HC, Song JW, Chong ST, O'Guinn ML, Lee JS, Lee IY, Park JH, Chae JS.** Detection of *Bartonella* species from ticks, mites and small mammals in Korea. *J Vet Sci* 2005, **6**, 327-334.
 17. **Kim CM, Kim MS, Park MS, Park JH, Chae JS.** Identification of *Ehrlichia chaffeensis*, *Anaplasma phagocytophilum*, and *A. bovis* in *Haemaphysalis longicornis* and *Ixodes persulcatus* ticks from Korea. *Vector Borne Zoonotic Dis* 2003, **3**, 17-26.
 18. **Kim CM, Yi YH, Yu DH, Lee MJ, Cho MR, Desai AR, Shringi S, Klein TA, Kim HC, Song JW, Baek LJ, Chong ST, O'Guinn ML, Lee JS, Lee IY, Park JH, Foley J, Chae JS.** Tick-borne rickettsial pathogens in ticks and small mammals in Korea. *Appl Environ Microbiol* 2006, **72**, 5766-5776.
 19. **Kim SY, Yun SM, Han MG, Lee IY, Lee NY, Jeong YE, Lee BC, Ju YR.** Isolation of tick-borne encephalitis viruses from wild rodents, South Korea. *Vector Borne Zoonotic Dis* 2008, **8**, 7-13.
 20. **La Scola B, Liang Z, Zeaiter Z, Houpijian P, Grimont PA, Raoult D.** Genotypic characteristics of two serotypes of *Bartonella henselae*. *J Clin Microbiol* 2002, **40**, 2002-2008.
 21. **Lee SO, Na DK, Kim CM, Li YH, Cho YH, Park JH, Lee JH, Eo SK, Klein TA, Chae JS.** Identification and prevalence of *Ehrlichia chaffeensis* infection in *Haemaphysalis longicornis* ticks from Korea by PCR, sequencing and phylogenetic analysis based on 16S rRNA gene. *J Vet Sci* 2005, **6**, 151-155.
 22. **Mathew JS, Ewing SA, Murphy GL, Kocan KM, Corstvet RE, Fox JC.** Characterization of a new isolate of *Ehrlichia platys* (order rickettsiales) using electron microscopy and polymerase chain reaction. *Vet Parasitol* 1997, **68**, 1-10.
 23. **Murphy GL, Ewing SA, Whitworth LC, Fox JC, Kocan AA.** A molecular and serologic survey of *Ehrlichia canis*, *E. chaffeensis*, and *E. ewingii* in dogs and ticks from Oklahoma. *Vet Parasitol* 1998, **79**, 325-339.
 24. **Norman AF, Regnery R, Jameson P, Greene C, Krause DC.** Differentiation of *Bartonella*-like isolates at the species level by PCR-restriction fragment length polymorphism in the citrate synthase gene. *J Clin Microbiol* 1995, **33**, 1797-1803.
 25. **Pahl A, Kühlbrandt U, Brune K, Röllinghoff M, Gessner A.** Quantitative detection of *Borrelia burgdorferi* by real-time PCR. *J Clin Microbiol* 1999, **37**, 1958-1963.
 26. **Park JH, Heo EJ, Choi KS, Dumler JS, Chae JS.** Detection of antibodies to *Anaplasma phagocytophilum* and *Ehrlichia chaffeensis* antigens in sera of Korean patients by western immunoblotting and indirect immunofluorescence assays. *Clin Diagn Lab Immunol* 2003, **10**, 1059-1064.
 27. **Park JW, Klein TA, Lee HC, Pacha LA, Ryu SH, Yeom JS, Moon SH, Kim TS, Chai JY, Oh MD, Choe KW.** Vivax malaria: a continuing health threat to the Republic of Korea. *Am J Trop Med Hyg* 2003, **69**, 159-167.
 28. **Park KH, Chang WH, Schwan TG.** Identification and characterization of Lyme disease spirochetes, *Borrelia burgdorferi* sensu lato, isolated in Korea. *J Clin Microbiol* 1993, **31**, 1831-1837.
 29. **Parola P, Davoust B, Raoult D.** Tick- and flea-borne rickettsial emerging zoonoses. *Vet Res* 2005, **36**, 469-492.
 30. **Regnery RL, Spruill CL, Plikaytis BD.** Genotypic identification of rickettsiae and estimation of intraspecies sequence divergence for portions of two rickettsial genes. *J Bacteriol* 1991, **173**, 1576-1589.
 31. **Schouls LM, Van De Pol I, Rijpkema SGT, Schot CS.** Detection and identification of *Ehrlichia*, *Borrelia burgdorferi* sensu lato, and *Bartonella* species in Dutch *Ixodes ricinus*

- ticks. J Clin Microbiol 1999, **37**, 2215-2222.
32. **Yoon SY, Gye MC, Lee HS.** Mammalian fauna in DMZ area. Korean J Environ Biol 2007, **25**, 215-222.
33. **Wang IN, Dykhuizen DE, Qiu W, Dunn JJ, Bosler EM, Luft BJ.** Genetic diversity of ospC in a local population of *Borrelia burgdorferi* sensu stricto. Genetics 1999, **151**, 15-30.